

---

**SUPPLEMENTAL LABELING**

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1571A-S1 0820**

<b>OXATHIPIPROLIN</b>	<b>GROUP</b>	<b>49</b>	<b>FUNGICIDE</b>
-----------------------	--------------	-----------	------------------

**Orondis®**

**Fungicide**

**This supplemental label expires on 08/17/2023 and must not be used or distributed after this date.**

Active Ingredient:

Oxathiapiprolin\*: ..... 18.7%

Other Ingredients: ..... 81.3%

Total:

100.0%

\* CAS No. 1003318-67-9

Orondis® is formulated as a suspension concentrate and contains 1.67 pounds of oxathiapiprolin per gallon of product.

**KEEP OUT OF REACH OF CHILDREN.  
CAUTION/PRECAUCIÓN**

**EPA Reg. No. 100-1571**

**All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using Orondis as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**syngenta®**

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### CROP USE DIRECTIONS

#### Nut, Tree Group 14-12

Crops (including all cultivars, varieties, and/or hybrids of these)			
African nut-tree	Coconut	Okari Nut	
Almond	Caquito Nut	Pachira Nut	
Beechnut	Dika Nut	Peach Palm Nut	
Brazil Nut	Ginkgo	Pecan	
Brazilian pine	Guiana chestnut	Pequi	
Bunya	Hazelnut (Filbert)	Pili Nut	
Bur Oak	Heartnut	Pine Nut	
Butternut	Hickory Nut	Pistachio	
Cajou Nut	Japanese horse-chestnut	Sapucaia Nut	
Candlenut	Macadamia Nut	Tropical Almond	
Cashew	Mongongo Nut	Walnut, black	
Chestnut	Monkey-Pot	Walnut, English	
Chinquapin	Monkey Puzzle Nut	Yellowhorn	
Target Disease	Rate fl oz/A (lb ai/A)	Application Timing	Use Directions
Phytophthora root and crown rot ( <i>Phytophthora</i> spp.)	4.8 – 9.6 (0.06 - 0.12)	<p>For <b>Resets or New Plantings</b>, make the first soil application at planting and up to one additional application 1-6 months later, coinciding with a root growth flush.</p> <p>For <b>Established Plantings</b>, make up to two soil applications at a 1- to 6-month interval, coinciding with root growth flush.</p>	<p>Apply as a drench or soil spray under the canopy around the base of the tree to the zone of maximum root density, or through irrigation water (micro-sprinkler or drip).</p> <p>For effective disease control, ensure that the product solution thoroughly wets the target root zone. If the application does not move the product to the root zone, and rain is not imminent, then follow with irrigation.</p>
<p><b>Resistance Management:</b></p> <ul style="list-style-type: none"> <li>• Refer to <b>Section 3.2</b>.</li> <li>• Make no more than 2 sequential applications of Orondis (or any other FRAC 49-containing product) before rotating to a fungicide with a different mode of action.</li> </ul>			
USE RESTRICTIONS			
<ol style="list-style-type: none"> <li>1) Refer to <b>Section 6.1</b> for additional product use restrictions.</li> <li>2) <b>Maximum Single Application Rate:</b> <b>DO NOT</b> exceed the maximum rate listed in the table for this application type.</li> <li>3) <b>Maximum Number of Applications per Year:</b> <b>DO NOT</b> make more than 2 applications at the maximum rate per year.</li> <li>4) <b>Minimum Application Interval:</b> 30 days</li> </ol>			

- 5) **Maximum Annual Rate:** 19.2 fl oz/A/year (0.25 lb ai/A/year)
  - a. **DO NOT** apply more than 0.25 lb ai/A/year of oxathiapiprolin-containing products.
- 6) **DO NOT** use in nursery production of transplanted crops.
- 7) **Pre-harvest Interval (PHI):** 30 days

## Tropical and Subtropical Fruit, Medium to Large Fruit, Smooth, Inedible Peel, Crop Subgroup 24B

Crops (including all cultivars, varieties, and/or hybrids of these)			
Abiu	Cupuacu	Mangosteen	Poshte
Akee apple	Etambe	Paho	Quandong
Avocado	Jatoba	Papaya	Sapote, black
Avocado, Guatemalan	Kei apple	Pawpaw, common	Sapote, green
Avocado, Mexican	Langsat	Pelipisan	Sapote, white
Avocado, West Indian	Lanjut	Pequi	Sataw
Bacury	Lucuma	Pequia	Screw-pine
Banana	Mabolo	Persimmon, American	Star apple
Banana, dwarf	Mango	Plantain	Tamarind-of-the-Indies
Binjal	Mango, horse	Pomegranate	Wild Loquat
Canistel	Mango, Saipan		
Target Disease	Rate fl oz/A (lb ai/A)	Application Timing	Use Directions
Phytophthora root and crown rot ( <i>Phytophthora</i> spp.)	4.8 – 9.6 (0.06 - 0.12)	<p><b>Soil Application:</b></p> <p>For <b>Resets or New Plantings</b>, make the first soil application at planting. An additional application may be made 30 days later if needed, coinciding with a root growth flush.</p> <p>For <b>Established Plantings</b>, make up to two soil applications on a 30 day interval, coinciding with root growth flush.</p>	<p>Apply as a drench or soil spray under the canopy around the base of the tree to the zone of maximum root density, or through irrigation water (micro-sprinkler or drip).</p> <p>For effective disease control, ensure that the product solution thoroughly wets the target root zone. If the application does not move the product to the root zone, and rain is not imminent, then follow with irrigation.</p>
<p><b>Resistance Management:</b></p> <ul style="list-style-type: none"> <li>• Refer to <b>Section 3.2</b>.</li> <li>• Make no more than 2 sequential applications of Orondis (or any other FRAC 49-containing product) before rotating to a fungicide with a different mode of action.</li> <li>• Do not follow soil applications of Orondis products with foliar applications of Orondis (or any other FRAC 49-containing product). Use either soil applications or foliar applications but not both for disease control.</li> </ul>			
USE RESTRICTIONS			
<ol style="list-style-type: none"> <li>1) Refer to <b>Section 6.1</b> for additional product use restrictions.</li> <li>2) <b>Maximum Single Application Rate:</b> <ol style="list-style-type: none"> <li>a. <b>Soil Applications:</b> <b>DO NOT</b> exceed maximum rate listed in the table for this application type.</li> </ol> </li> <li>3) <b>Maximum Number of Applications per Year:</b></li> </ol>			

- a. **Soil applications: DO NOT** make more than 2 applications at the maximum rate per year.
- 4) **Minimum Application Interval:** 30 days soil
- 5) **Maximum Annual Rate:**
  - a. **Soil Applications:** 19.2 fl oz/A/year (0.25 lb ai/A/year)
  - b. **DO NOT** apply more than 0.25 lb ai/A/year of soil-applied oxathiapiprolin-containing products.
- 6) **DO NOT** use in nursery production of transplanted crops.
- 7) **Pre-harvest Interval (PHI):**
  - a. **Soil Applications:** 30 days

Orondis® and the Syngenta logo are trademarks of a Syngenta Group Company

©2020 Syngenta

Chron :

000100-01571.20190322E.ORONDIS\_SUPP\_0319-CL

SCP 1571A-S1 0820 – kdy 9/1/20